Supporting California Standards

Mathematics Continuum: Grades 6-8



Standards for Mathematical Practice (K-12)

- 1. Make sense of problems and persevere in solving them.
- 2. Reason abstractly and quantitatively.
- 3. Construct viable arguments and critique the reasoning of others
- 4. Model with mathematics

- 5. Use appropriate tools strategically
- 6. Attend to precision.
- 7. Look for and make use of structure.
- 8. Look for and express regularity in repeated reasoning.

Ratios and Proportional Relationships			
Grade 6	Grade 7	Grade 8	
 Understand ratio concepts and use ratio reasoning to solve problems. 6.RP.1. Understand the concept of a ratio and use ratio language to describe a ratio relationship between two quantities. For example, "The ratio of wings to beaks in the bird house at the zoo was 2:1, because for every 2 wings there was 1 beak." "For every vote candidate A received, candidate C received nearly three votes." 6.RP.2. Understand the concept of a unit rate a/b associated with a ratio a:b with b ≠ 0, and use rate language in the context of a ratio relationship. For example, "This recipe has a ratio of 3 cups of flour to 4 cups of sugar, so there is 3/4 cup of flour for each cup of sugar." "We paid \$75 for 15 hamburgers, which is a rate of \$5 per hamburger."¹ [¹ Expectations for unit rates in this grade are limited to noncomplex fractions.] 6.RP.3. Use ratio and rate reasoning to solve real-world and mathematical problems, e.g., by reasoning about tables of equivalent ratios, tape diagrams, double number line diagrams, or equations. a. Make tables of equivalent ratios relating quantities with whole number measurements, find missing values in the tables, and plot the pairs of values on the coordinate plane. Use tables to compare ratios. b. Solve unit rate problems including those involving unit pricing and constant speed. For example, if it took 7 hours to mow 4 lawns, then at that rate, how many lawns could be mowed in 35 hours? At what rate were lawns being mowed? c. Find a percent of a quantity as a rate per 100 (e.g., 30% of a quantity means 30/100 times the quantity); solve problems involving finding the whole, given a part and the percent. d. Use ratio reasoning to convert measurement units; manipulate and transform units appropriately when multiplying or dividing quantities. 	Analyze proportional relationships and use them to solve real-world and mathematical problems. 7.RP.1. Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour. 7.RP.2. Recognize and represent proportional relationships between quantities. a. Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin. b. Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships. c. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn. d. Explain what a point (x, y) on the graph of a proportional relationship means in terms of the situation, with special attention to the points (0, 0) and (1, r) where r is the unit rate. 7.RP.3. Use proportional relationships to solve multistep ratio and percent problems. Examples: simple interest, tax, markups and markdowns, gratuities and commissions, fees, percent increase and decrease, percent error.	See 8.EE.5	



The Number System			
Grade 6	Grade 7	Grade 8	
Apply and extend previous understandings of multiplication and division to divide fractions by fractions.	Apply and extend previous understandings of operations with fractions to	Know that there are	
6.NS.1. Interpret and compute quotients of fractions, and solve word problems involving division of fractions by fractions, e.g., by using	add, subtract, multiply, and divide rational numbers.	numbers that are not	
visual fraction models and equations to represent the problem. For example, create a story context for (2/3) ÷ (3/4) and use a	7.NS.1. Apply and extend previous understandings of addition and subtraction to	rational, and approximate	
visual fraction model to show the quotient; use the relationship between multiplication and division to explain that $(2/3) \div (3/4) =$	add and subtract rational numbers; represent addition and subtraction on	them by rational numbers.	
8/9 because 3/4 of 8/9 is 2/3. (In general, $(a/b) \div (c/d) = ad/bc$.) How much chocolate will each person get if 3 people share 1/2 lb	a horizontal or vertical number line diagram.	8.NS.1. Know that numbers	
of chocolate equally? How many 3/4-cup servings are in 2/3 of a cup of yogurt? How wide is a rectangular strip of land with	a. Describe situations in which opposite quantities combine to make 0. For	that are not rational	
length 3/4 mi and area 1/2 square mi?	example, a hydrogen atom has 0 charge because its two constituents	are called irrational.	
Compute fluently with multi-digit numbers and find common factors and multiples.	are oppositely charged.	Understand informally	
6.NS.2. Fluently divide multi-digit numbers using the standard algorithm.	b. Understand $p + q$ as the number located a distance $ q $ from p , in the	that every number has	
6.NS.3. Fluently add, subtract, multiply, and divide multi-digit decimals using the standard algorithm for each operation.	positive or negative direction depending on whether q is positive or	a decimal expansion;	
6.NS.4. Find the greatest common factor of two whole numbers less than or equal to 100 and the least common multiple of two whole	negative. Show that a number and its opposite have a sum of 0 (are	for rational numbers	
numbers less than or equal to 12. Use the distributive property to express a sum of two whole numbers 1–100 with a common	additive inverses). Interpret sums of rational numbers by describing real-	show that the decimal	
factor as a multiple of a sum of two whole numbers with no common factor. For example, express 36 + 8 as 4 (9 + 2).	world contexts.	expansion repeats	
Apply and extend previous understandings of numbers to the system of rational numbers.	c. Understand subtraction of rational numbers as adding the additive	eventually, and	
6.NS.5. Understand that positive and negative numbers are used together to describe quantities having opposite directions or values	inverse, $p - q = p + (-q)$. Show that the distance between two rational	convert a decimal	
(e.g., temperature above/below zero, elevation above/below sea level, credits/debits, positive/negative electric charge); use	numbers on the number line is the absolute value of their difference, and	expansion which	
positive and negative numbers to represent quantities in real-world contexts, explaining the meaning of 0 in each situation.	apply this principle in real-world contexts.	repeats eventually into	
6.NS.6. Understand a rational number as a point on the number line. Extend number line diagrams and coordinate axes familiar from	d. Apply properties of operations as strategies to add and subtract rational	a rational number. 8.NS.2. Use rational	
previous grades to represent points on the line and in the plane with negative number coordinates.	numbers. 7.NS.2. Apply and extend previous understandings of multiplication and division	approximations of	
a. Recognize opposite signs of numbers as indicating locations on opposite sides of 0 on the number line; recognize that the opposite of the opposite of a number is the number itself, e.g., -(-3) = 3, and that 0 is its own opposite.	and of fractions to multiply and divide rational numbers.	irrational numbers to	
b. Understand signs of numbers in ordered pairs as indicating locations in quadrants of the coordinate plane; recognize that when	a. Understand that multiplication is extended from fractions to rational	compare the size of	
two ordered pairs differ only by signs, the locations of the points are related by reflections across one or both axes.	numbers by requiring that operations continue to satisfy the properties of	irrational numbers,	
c. Find and position integers and other rational numbers on a horizontal or vertical number line diagram; find and position pairs of	operations, particularly the distributive property, leading to products such	locate them	
integers and other rational numbers on a coordinate plane.	as $(-1)(-1) = 1$ and the rules for multiplying signed numbers. Interpret	approximately on a	
6.NS.7. Understand ordering and absolute value of rational numbers.	products of rational numbers by describing real-world contexts.	number line diagram,	
a. Interpret statements of inequality as statements about the relative position of two numbers on a number line diagram. <i>For</i>	b. Understand that integers can be divided, provided that the divisor is not	and estimate the value	
example, interpret $-3 > -7$ as a statement that -3 is located to the right of -7 on a number line oriented from left to right.	zero, and every quotient of integers (with non-zero divisor) is a rational	of expressions (e.g.,	
b. Write, interpret, and explain statements of order for rational numbers in real-world contexts. For example, write $-3^{\circ}C > -7^{\circ}C$ to	number. If p and q are integers, then $-(p/q) = (-p)/q = p/(-q)$. Interpret	π2). For example, by	
express the fact that –3°C is warmer than –7°C.	quotients of rational numbers by describing real world contexts.	truncating the decimal	
c. Understand the absolute value of a rational number as its distance from 0 on the number line; interpret absolute value as	c. Apply properties of operations as strategies to multiply and divide rational	expansion of √2, show	
magnitude for a positive or negative quantity in a real-world situation. For example, for an account balance of –30 dollars, write	numbers.	that √2 is between 1	
-30 = 30 to describe the size of the debt in dollars.	d. Convert a rational number to a decimal using long division; know that the	and 2, then between	
d. Distinguish comparisons of absolute value from statements about order. For example, recognize that an account balance less	decimal form of a rational number terminates in 0s or eventually repeats.	1.4 and 1.5, and	
than –30 dollars represents a debt greater than 30 dollars.	7.NS.3. Solve real-world and mathematical problems involving the four	explain how to	
6.NS.8. Solve real-world and mathematical problems by graphing points in all four quadrants of the coordinate plane. Include use of	operations with rational numbers. [Computations with rational numbers	continue on to get	
coordinates and absolute value to find distances between points with the same first coordinate or the same second coordinate.	extend the rules for manipulating fractions to complex fractions.]	better approximations.	



Expressions and Equations			
Grade 6	Grade 7	Grade 8	
Apply and extend previous understandings of arithmetic to algebraic expressions.	Use properties of operations to generate equivalent expressions.	Work with radicals and integer exponents.	
6.EE.1. Write and evaluate numerical expressions involving whole-number exponents.	7.EE.1. Apply properties of operations as strategies to add, subtract,	8.EE.1. Know and apply the properties of integer exponents to generate	
6.EE.2. Write, read, and evaluate expressions in which letters stand for numbers.	factor, and expand linear expressions with rational coefficients.	equivalent numerical expressions. For example, $3^2 \times 3^{-5} = 3^{-3} = 1/3^3$	
 a. Write expressions that record operations with numbers and with letters 	7.EE.2. Understand that rewriting an expression in different forms in a	1/27.	
standing for numbers. For example, express the calculation "Subtract y from	problem context can shed light on the problem and how the	8.EE.2. Use square root and cube root symbols to represent solutions to	
5" as 5 − y.	quantities in it are related. For example, a + 0.05a = 1.05a means	equations of the form $x^2 = p$ and $x^3 = p$, where p is a positive rational	
b. Identify parts of an expression using mathematical terms (sum, term, product,	that "increase by 5%" is the same as "multiply by 1.05."	number. Evaluate square roots of small perfect squares and cube roots	
factor, quotient, coefficient); view one or more parts of an expression as a	Solve real-life and mathematical problems using numerical and	of small perfect cubes. Know that $\sqrt{2}$ is irrational.	
single entity. For example, describe the expression 2 (8 + 7) as a product of	algebraic expressions and equations.	8.EE.3. Use numbers expressed in the form of a single digit times an integer	
two factors; view (8 + 7) as both a single entity and a sum of two terms.	7.EE.3. Solve multi-step real-life and mathematical problems posed with	power of 10 to estimate very large or very small quantities, and to	
c. Evaluate expressions at specific values of their variables. Include expressions	positive and negative rational numbers in any form (whole	express how many times as much one is than the other. For example,	
that arise from formulas used in real-world problems. Perform arithmetic	numbers, fractions, and decimals), using tools strategically. Apply	estimate the population of the United States as 3 × 10 ⁸ and the	
operations, including those involving whole-number exponents, in the	properties of operations to calculate with numbers in any form;	population of the world as 7×10^9 , and determine that the world	
conventional order when there are no parentheses to specify a particular	convert between forms as appropriate; and assess the	population is more than 20 times larger.	
order (Order of Operations). For example, use the formulas $V = s^3$ and $A = 6$	reasonableness of answers using mental computation and	8.EE.4. Perform operations with numbers expressed in scientific notation,	
s^2 to find the volume and surface area of a cube with sides of length $s = 1/2$.	estimation strategies. For example: If a woman making \$25 an	including problems where both decimal and scientific notation are used.	
6.EE.3. Apply the properties of operations to generate equivalent expressions. <i>For</i>	hour gets a 10% raise, she will make an additional 1/10 of her	Use scientific notation and choose units of appropriate size for	
example, apply the distributive property to the expression $3(2 + x)$ to produce the	salary an hour, or \$2.50, for a new salary of \$27.50. If you want to	measurements of very large or very small quantities (e.g., use millimeters	
equivalent expression 6 + 3x; apply the distributive property to the expression 24x	place a towel bar 9 3/4 inches long in the center of a door that is	per year for seafloor spreading). Interpret scientific notation that has been	
+ 18y to produce the equivalent expression 6 (4x + 3y); apply properties of	27 1/2 inches wide, you will need to place the bar about 9 inches	generated by technology.	
operations to $y + y + y$ to produce the equivalent expression 3y.	from each edge; this estimate can be used as a check on the exact	Understand the connections between proportional relationships, lines, and	
6.EE.4. Identify when two expressions are equivalent (i.e., when the two expressions	computation.	linear equations.	
name the same number regardless of which value is substituted into them). For	7.EE.4. Use variables to represent quantities in a real-world or	8.EE.5. Graph proportional relationships, interpreting the unit rate as the slope of	
example, the expressions $y + y + y$ and $3y$ are equivalent because they name the	mathematical problem, and construct simple equations and	the graph. Compare two different proportional relationships represented	
same number regardless of which number y stands for.	inequalities to solve problems by reasoning about the quantities.	in different ways. For example, compare a distance-time graph to a	
Reason about and solve one-variable equations and inequalities.	a. Solve word problems leading to equations of the form $px + q = \frac{1}{2} \left(\frac{1}{2} + \frac{1}{2} \right) \left(\frac{1}{2} + \frac{1}{2} \right)$	distance-time equation to determine which of two moving objects has	
6.EE.5. Understand solving an equation or inequality as a process of answering a	r and $p(x + q) = r$, where p , q , and r are specific rational	greater speed.	
question: which values from a specified set, if any, make the equation or inequality	numbers. Solve equations of these forms fluently. Compare an	8.EE.6. Use similar triangles to explain why the slope <i>m</i> is the same between any two distinct points on a non-vertical line in the coordinate plane; derive	
true? Use substitution to determine whether a given number in a specified set	algebraic solution to an arithmetic solution, identifying the	1	
makes an equation or inequality true. 6.EE.6. Use variables to represent numbers and write expressions when solving a real-	sequence of the operations used in each approach. For	the equation $y = mx$ for a line through the origin and the equation $y = mx$ + b for a line intercepting the vertical axis at b .	
world or mathematical problem; understand that a variable can represent an	example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?		
unknown number, or, depending on the purpose at hand, any number in a	b. Solve word problems leading to inequalities of the form px + q	Analyze and solve linear equations and pairs of simultaneous linear	
specified set.	> r or px + q < r, where p, q, and r are specific rational	equations. 8.EE.7. Solve linear equations in one variable.	
6.EE.7. Solve real-world and mathematical problems by writing and solving equations of	numbers. Graph the solution set of the inequality and interpret	a. Give examples of linear equations in one variable with one solution,	
the form $x + p = q$ and $px = q$ for cases in which p , q and x are all nonnegative	it in the context of the problem. For example: As a salesperson,	infinitely many solutions, or no solutions. Show which of these	
rational numbers.	you are paid \$50 per week plus \$3 per sale. This week you	possibilities is the case by successively transforming the given	
6.EE.8. Write an inequality of the form $x > c$ or $x < c$ to represent a constraint or condition	want your pay to be at least \$100. Write an inequality for the	equation into simpler forms, until an equivalent equation of the form <i>x</i> =	
in a real-world or mathematical problem. Recognize that inequalities of the form <i>x</i>	number of sales you need to make, and describe the solutions.	a, $a = a$, or $a = b$ results (where a and b are different numbers).	
in a roal-world of mathematical problem. Neoognize that inequalities of the form X	Transport of sales you need to make, and describe the solutions.	α , $\alpha = \alpha$, or $\alpha = \nu$ results (where α and ν are different numbers).	



> c or x < c have infinitely many solutions; represent solutions of such inequalities on number line diagrams.

Represent and analyze quantitative relationships between dependent and independent variables.

6.EE.9. Use variables to represent two quantities in a real-world problem that change in relationship to one another; write an equation to express one quantity, thought of as the dependent variable, in terms of the other quantity, thought of as the independent variable. Analyze the relationship between the dependent and independent variables using graphs and tables, and relate these to the equation. For example, in a problem involving motion at constant speed, list and graph ordered pairs of distances and times, and write the equation d = 65t to represent the relationship between distance and time.

 Solve linear equations with rational number coefficients, including equations whose solutions require expanding expressions using the distributive property and collecting like terms.

8.EE.8. Analyze and solve pairs of simultaneous linear equations.

- a. Understand that solutions to a system of two linear equations in two variables correspond to points of intersection of their graphs, because points of intersection satisfy both equations simultaneously.
- b. Solve systems of two linear equations in two variables algebraically, and estimate solutions by graphing the equations. Solve simple cases by inspection. For example, 3x + 2y = 5 and 3x + 2y = 6 have no solution because 3x + 2y cannot simultaneously be 5 and 6.
- c. Solve real-world and mathematical problems leading to two linear equations in two variables. For example, given coordinates for two pairs of points, determine whether the line through the first pair of points intersects the line through the second pair.

Functions Function Functions Functions Functions Functions Function Fun		
Grade 6	Grade 7	Grade 8
None	None	 Define, evaluate, and compare functions. 8.F.1. Understand that a function is a rule that assigns to each input exactly one output. The graph of a function is the set of ordered pairs consisting of an input and the corresponding output. [Function notation is not required in Grade 8.] 8.F.2. Compare properties of two functions each represented in a different way (algebraically, graphically, numerically in tables, or by verbal descriptions). For example, given a linear function represented by a table of values and a linear function represented by an algebraic expression, determine which function has the greater rate of change. 8.F.3. Interpret the equation y = mx + b as defining a linear function, whose graph is a straight line; give examples of functions that are not linear. For example, the function A = s² giving the area of a square as a function of its side length is not linear because its graph contains the points (1,1), (2,4) and (3,9), which are not on a straight line. Use functions to model relationships between quantities. 8.F.4. Construct a function to model a linear relationship between two quantities. Determine the rate of change and initial value of the function from a description of a relationship or from two (x, y) values, including reading these from a table or from a graph. Interpret the rate of change and initial value of a linear function in terms of the situation it models, and in terms of its graph or a table of values. 8.F.5. Describe qualitatively the functional relationship between two quantities by analyzing a graph (e.g., where the function is increasing or decreasing, linear or nonlinear). Sketch a graph that exhibits the qualitative features of a function that has been described verbally.



Geometry			
Grade 6	Grade 7	Grade 8	
Solve real-world and mathematical problems	Draw, construct, and describe geometrical figures and	Understand congruence and similarity using physical models, transparencies, or geometry	
involving area, surface area, and volume.	describe the relationships between them.	software.	
6.G.1. Find the area of right triangles, other triangles,	7.G.1. Solve problems involving scale drawings of geometric	8.G.1. Verify experimentally the properties of rotations, reflections, and translations:	
special quadrilaterals, and polygons by composing	figures, including computing actual lengths and areas	a. Lines are taken to lines, and line segments to line segments of the same length.	
into rectangles or decomposing into triangles and	from a scale drawing and reproducing a scale drawing	b. Angles are taken to angles of the same measure.	
other shapes; apply these techniques in the	at a different scale.	c. Parallel lines are taken to parallel lines.	
context of solving real-world and mathematical	7.G.2. Draw (freehand, with ruler and protractor, and with	8.G.2. Understand that a two-dimensional figure is congruent to another if the second can be obtained	
problems.	technology) geometric shapes with given conditions.	from the first by a sequence of rotations, reflections, and translations; given two congruent figures,	
6.G.2. Find the volume of a right rectangular prism with	Focus on constructing triangles from three measures	describe a sequence that exhibits the congruence between them.	
fractional edge lengths by packing it with unit	of angles or sides, noticing when the conditions	8.G.3. Describe the effect of dilations, translations, rotations, and reflections on two-dimensional figures	
cubes of the appropriate unit fraction edge lengths,	determine a unique triangle, more than one triangle, or	using coordinates.	
and show that the volume is the same as would be	no triangle.	8.G.4. Understand that a two-dimensional figure is similar to another if the second can be obtained from	
found by multiplying the edge lengths of the prism.	7.G.3. Describe the two-dimensional figures that result from	the first by a sequence of rotations, reflections, translations, and dilations; given two similar two-	
Apply the formulas $V = l w h$ and $V = b h$ to find	slicing three-dimensional figures, as in plane sections	dimensional figures, describe a sequence that exhibits the similarity between them.	
volumes of right rectangular prisms with fractional	of right rectangular prisms and right rectangular	8.G.5. Use informal arguments to establish facts about the angle sum and exterior angle of triangles,	
edge lengths in the context of solving real-world	pyramids.	about the angles created when parallel lines are cut by a transversal, and the angle-angle criterion	
and mathematical problems.	Solve real-life and mathematical problems involving angle	for similarity of triangles. For example, arrange three copies of the same triangle so that the sum of	
6.G.3. Draw polygons in the coordinate plane given	measure, area, surface area, and volume.	the three angles appears to form a line, and give an argument in terms of transversals why this is	
coordinates for the vertices; use coordinates to	7.G.4. Know the formulas for the area and circumference of a	SO.	
find the length of a side joining points with the	circle and use them to solve problems; give an	Understand and apply the Pythagorean Theorem.	
same first coordinate or the same second	informal derivation of the relationship between the circumference and area of a circle.	8.G.6. Explain a proof of the Pythagorean Theorem and its converse.	
coordinate. Apply these techniques in the context		8.G.7. Apply the Pythagorean Theorem to determine unknown side lengths in right triangles in real-world	
of solving real-world and mathematical problems. 6.G.4. Represent three-dimensional figures using nets	7.G.5. Use facts about supplementary, complementary,	and mathematical problems in two and three dimensions. 8.G.8. Apply the Pythagorean Theorem to find the distance between two points in a coordinate system.	
made up of rectangles and triangles, and use the	vertical, and adjacent angles in a multi-step problem to write and solve simple equations for an unknown angle	6.6.6. Apply the Fythagorean Theorem to lind the distance between two points in a coordinate system.	
nets to find the surface area of these figures. Apply	in a figure.	Solve real-world and mathematical problems involving volume of cylinders, cones, and spheres.	
these techniques in the context of solving real-	7.G.6. Solve real-world and mathematical problems involving	8.G.9. Know the formulas for the volumes of cones, cylinders, and spheres and use them to solve real-	
world and mathematical problems.	area, volume and surface area of two- and three-	world and mathematical problems.	
world and mathematical problems.	dimensional objects composed of triangles,	world and mathematical problems.	
	quadrilaterals, polygons, cubes, and right prisms.		
	quadriaterate, perjagone, eases, and right phonie.		

Supporting California Standards

Mathematics Continuum: Grades 6-8



Statistics and Probability			
Grade 6	Grade 7	Grade 8	
Develop understanding of statistical variability.	Use random sampling to draw inferences about a population.	Investigate patterns of association in bivariate	
6.SP.1. Recognize a statistical question as one that	7.SP.1. Understand that statistics can be used to gain information about a population by examining a sample of the population;	data.	
anticipates variability in the data related to the	generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that	8.SP.1. Construct and interpret scatter plots for	
question and accounts for it in the answers. For	random sampling tends to produce representative samples and support valid inferences.	bivariate measurement data to investigate	
example, "How old am I?" is not a statistical	7.SP.2. Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate	patterns of association between two	
question, but "How old are the students in my	multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate	quantities. Describe patterns such as	
school?" is a statistical question because one	the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on	clustering, outliers, positive or negative	
anticipates variability in students' ages.	randomly sampled survey data. Gauge how far off the estimate or prediction might be.	association, linear association, and nonlinear	
6.SP.2. Understand that a set of data collected to	Draw informal comparative inferences about two populations.	association.	
answer a statistical question has a distribution	7.SP.3. Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the	8.SP.2. Know that straight lines are widely used to	
which can be described by its center, spread, and	difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on	model relationships between two quantitative	
overall shape.	the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability (mean absolute	variables. For scatter plots that suggest a	
6.SP.3. Recognize that a measure of center for a	deviation) on either team; on a dot plot, the separation between the two distributions of heights is noticeable.	linear association, informally fit a straight line,	
numerical data set summarizes all of its values	7.SP.4. Use measures of center and measures of variability for numerical data from random samples to draw informal comparative	and informally assess the model fit by judging	
with a single number, while a measure of variation	inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are	the closeness of the data points to the line.	
describes how its values vary with a single	generally longer than the words in a chapter of a fourth-grade science book.	8.SP.3. Use the equation of a linear model to solve	
number.	Investigate chance processes and develop, use, and evaluate probability models.	problems in the context of bivariate	
Summarize and describe distributions.	7.SP.5. Understand that the probability of a chance event is a number between 0 and 1 that expresses the likelihood of the event	measurement data, interpreting the slope and	
6.SP.4. Display numerical data in plots on a number	occurring. Larger numbers indicate greater likelihood. A probability near 0 indicates an unlikely event, a probability around 1/2	intercept. For example, in a linear model for a	
line, including dot plots, histograms, and box	indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event.	biology experiment, interpret a slope of 1.5	
plots.	7.SP.6. Approximate the probability of a chance event by collecting data on the chance process that produces it and observing its long-	cm/hr as meaning that an additional hour of	
6.SP.5. Summarize numerical data sets in relation to	run relative frequency, and predict the approximate relative frequency given the probability. For example, when rolling a number	sunlight each day is associated with an	
their context, such as by:	cube 600 times, predict that a 3 or 6 would be rolled roughly 200 times, but probably not exactly 200 times.	additional 1.5 cm in mature plant height.	
 Reporting the number of observations. 	7.SP.7. Develop a probability model and use it to find probabilities of events. Compare probabilities from a model to observed	8.SP.4. Understand that patterns of association	
b. Describing the nature of the attribute under	frequencies; if the agreement is not good, explain possible sources of the discrepancy.	can also be seen in bivariate categorical data	
investigation, including how it was measured	a. Develop a uniform probability model by assigning equal probability to all outcomes, and use the model to determine probabilities	by displaying frequencies and relative	
and its units of measurement.	of events. For example, if a student is selected at random from a class, find the probability that Jane will be selected and the	frequencies in a two-way table. Construct	
c. Giving quantitative measures of center	probability that a girl will be selected.	and interpret a two-way table summarizing	
(median and/or mean) and variability	b. Develop a probability model (which may not be uniform) by observing frequencies in data generated from a chance process. For	data on two categorical variables collected	
(interquartile range and/or mean absolute	example, find the approximate probability that a spinning penny will land heads up or that a tossed paper cup will land open-end	from the same subjects. Use relative	
deviation), as well as describing any overall	down. Do the outcomes for the spinning penny appear to be equally likely based on the observed frequencies?	frequencies calculated for rows or columns to	
pattern and any striking deviations from the	7.SP.8. Find probabilities of compound events using organized lists, tables, tree diagrams, and simulation.	describe possible association between the	
overall pattern with reference to the context in	a. Understand that, just as with simple events, the probability of a compound event is the fraction of outcomes in the sample space	two variables. For example, collect data from	
which the data were gathered.	for which the compound event occurs.	students in your class on whether or not they	
d. Relating the choice of measures of center and	b. Represent sample spaces for compound events using methods such as organized lists, tables and tree diagrams. For an event	have a curfew on school nights and whether	
variability to the shape of the data distribution	described in everyday language (e.g., "rolling double sixes"), identify the outcomes in the sample space which compose the event.	or not they have assigned chores at home. Is	
and the context in which the data were	c. Design and use a simulation to generate frequencies for compound events. For example, use random digits as a simulation tool to	there evidence that those who have a curfew	
gathered.	approximate the answer to the question: If 40% of donors have type A blood, what is the probability that it will take at least 4	also tend to have chores?	
	donors to find one with type A blood?		

